

To appear in the ApJ, 10 October issue

The M/L_* ratio of young star clusters in galactic mergers

C. M. Boily, A. Lançon

Observatoire astronomique, 11 rue de l'Université, F-67000 Strasbourg, France;
{cmb,lancon}@astro.u-strasbg.fr

S. Deiters and D.C. Heggie

School of Mathematics, University of Edinburgh, King's Building, Edinburgh, Scotland;
{s.deiters,d.c.heggie}@ed.ac.uk

ABSTRACT

We point out a strong time-evolution of the mass-to-light conversion factor η commonly used to estimate masses of dense star clusters from observed cluster radii and stellar velocity dispersions. We use a gas-dynamical model coupled with the Cambridge stellar evolution tracks to compute line-of-sight velocity dispersions and half-light radii weighted by the luminosity. Stars at birth are assumed to follow the Salpeter mass function in the range $[0.15\text{--}17 M_\odot]$. We find that η , and hence the estimated cluster mass, increases by factors as large as 3 over time-scales of 20 million years. Increasing the upper mass limit to $50 M_\odot$ leads to a sharp rise of similar amplitude but in as little as 10 million years. Fitting truncated isothermal (Michie-King) models to the projected light profile leads to over-estimates of the concentration parameter c of $\delta c \approx 0.3$ compared to the same functional fit applied to the projected mass density.

Subject headings: stellar dynamics – stellar evolution – NGC4038 – M82 – R136
 – numerical method

1 Introduction

The formation of star clusters in bursts of star formation during galactic mergers has attracted much attention since the ground-breaking study by Schweizer (1986). Young clusters can account for $\sim 20\%$ of the UV light flux of their sample starburst galaxies (compared with $< 1\%$ for the Milky Way; e.g. Meurer et al. 1995). Such numbers bring to focus

the role that star formation in clusters plays in shaping the overall (galactic) stellar mass function. Proto-typical cases where cluster formation has been a spectacular manifestation of interaction-induced starbursts are the merging systems NGC 4038/39 (the Antennæ) and the nearby galaxy M82 (recent interaction with M81). Many of the brightest clusters in these galaxies have estimated ages on the order of 10^7 years, based on optical and near-IR spectra. High resolution spectroscopic data and HST images have been used to measure velocity dispersions and estimate virial masses. The line-of-sight velocity dispersion σ_{1d} (LOSVD) relates to the mass M and projected half-light radius r_{ph} of a cluster in virial equilibrium through

$$M = \eta \frac{r_{ph} \sigma_{1d}^2}{G} . \quad (1)$$

where η is a dimensionless free parameter. A number of authors have set $\eta \approx 10$ in their studies to derive M from (1) (Sternberg 1998; Mengel et al. 2002; Smith & Gallagher 2001; McCrady et al. 2003; Maraston et al. 2004). McCrady et al give a derivation of this value of η . Mengel et al. (2002) quote a range from 5.6 – 9.7 for King (1966) models with concentration parameter in the range 0.5 to 2.5, which corresponds to most galactic globular clusters. We discuss the value of η further below. Dynamical masses derived from (1) have been compared to the stellar masses of synthetic populations, using both standard (field) and non-standard stellar initial mass functions (henceforth IMF; see Kroupa 2002 for a review). The data were found to be inconsistent with a universal IMF (Mengel et al. 2002; Smith & Gallagher 2001). In particular, several clusters in M82 were found to be over-luminous with respect to their estimated mass (low mass-to-light ratio, M/L_*). This suggests that M82 clusters may form with a top-heavy stellar IMF (Smith & Gallagher 2001; McCrady et al. 2003).

These conclusions hinge on the precise value of η . The above studies have assumed no time-variation of η , based on the belief that the structural parameters of young clusters are unlikely to have changed since the time they were born. A further implicit assumption made when applying (1) is that the stellar subpopulations sampled are all equally representative of the dynamics as a whole. This simplification is normally justified on the grounds that the estimated ages of the clusters are too short to allow for gradients in their spatial distribution and kinematics to develop from internal evolution. However this train of thought stems from the derivation of a long relaxation time t_{rh} for single-population clusters. Expressed in terms of the dynamical time t_{cr} evaluated at the half-mass radius R_h (Meylan & Heggie 1997, §7)

$$\frac{t_{rh}}{t_{cr}} \simeq 0.138 \left[\frac{R_h}{R_g} \right]^{3/2} \frac{N}{\ln 0.4N} \quad (2)$$

where $R_g = GM^2/|W|$ is the gravitational radius and the ratio $R_h/R_g \approx$ unity for a wide number of model fits to observed clusters. With a spectrum of masses, the trend toward equipartition of kinetic energy speeds up evolution, and mass segregation now develops on a time-scale given by (Farouki & Salpeter 1982; Spitzer 1987)

$$\frac{t_{ms}}{t_{rh}} \approx \frac{\pi}{3} \frac{\langle m \rangle}{\max\{m\}} \frac{\bar{\rho}}{\rho} \left[\frac{R_h}{R_g} \right]^{3/2} \quad (3)$$

where $\bar{\rho} \equiv M/2/(4\pi R_h^3/3)$ is the mean density inside the half-mass radius (an over-line denotes averaging over space, and brackets averaging by mass). The numerical coefficients entering (3) are derived from a stellar IMF and mass distribution. The Galactic-field IMF covers a range from $\approx 0.08 M_\odot$ to $\gtrsim 60 M_\odot$ (O-stars) and possibly all the way to $\approx 100 M_\odot$ (Kroupa 2002). The mean mass for this IMF is $\langle m \rangle \approx 1.33 M_\odot$ and therefore the ratio $\langle m \rangle/\max\{m\}$ lies in the range $0.013 - 0.022$ for the chosen upper limits. This dramatically reduces the mass-segregation time-scale (3) and bears on the parameter η (through time-variation of r_{ph} and σ_{1d}) since the brightest, most massive member stars are also those that undergo the most significant segregation and inward migration. The purpose of this Letter is to show with numerical modelling that, through this process, the mass conversion factor η of massive clusters evolves by a factor of a few over periods as short as 20 millions years, contrary to expectations of zero evolution.

2 Numerical method : GasTel

The equations of motion were integrated numerically based on the gas-dynamical approach pioneered by Larson (1970) and developed further by Louis & Spurzem (1991) to include anisotropic velocity fields. The method leans on an analogy between exchange of kinetic energy through star-star interactions and the classical heat-diffusion process of fluid dynamics (Lynden-Bell & Eggleton 1980; Heggie & Ramamani 1989). The implementation in spherical coordinates we used is largely due to Louis & Spurzem (1991) and Giersz & Spurzem (1994) but extended to include a spectrum of stellar masses (Spurzem & Takahashi 1995).

The mass spectrum is sampled at constant logarithmic increments in the interval $\{m_0, m_1\}$; we used 14 bins in our standard runs ($\delta \ln m \approx 0.329$). Star-star interactions (including those between stars of the same mass bin) lead to the diffusion of kinetic energy. Roughly speaking, the resulting change in the velocity dispersion of each mass bin causes a readjustment of the density profile. This is obtained using a semi-implicit Henyey integration. The gravitational potential is then updated by applying Poisson's equation (see Louis & Spurzem 1991).

We checked that the mass-segregation time (3) obtained for GasTel is in quantitative agreement with N-body and Fokker-Planck integrators by computing the evolution of Plummer models with three species of stars. (cf. Spitzer & Shull 1975, Fig. 1). Spurzem & Takahashi (1995) report excellent agreement with (3) from their two-component test calculations.

2.1 Stellar evolution

The different mass components are evolved according to the Cambridge tracks (Pols et al. 1998; Hurley et al. 2000). The tracks are efficiently coded in the form of fitting functions of the kind first presented by Eggleton, Fitchett & Tout (1989). The functions return the current bolometric luminosity, radius, mass and metal abundance for given time and initial metal abundance z_o ; in this contribution we set $z_o = 0.02$ (solar abundance) throughout.

A filter can be applied to the bolometric luminosity from model stellar atmospheres, and the total flux in a specified waveband read from the Basel stellar library (Lejeune et al. 1998). We have mapped the stellar luminosity near the strongest near-IR CO bandhead ($\lambda \simeq 2.2 - 2.29\mu m$) and the CaII triplet ($\lambda \simeq 8200 - 8600\text{\AA}$).

3 Flux-weighted scheme & stellar IMF

Low gradients in metal abundances in young clusters can be interpreted to imply a spatial distribution of stars independent of their mass at the time the clusters formed (e.g. Suntzeff 1993). All the calculations presented here have no built-in segregation initially.

The flux-weighted LOSVD is obtained, first by averaging the square velocity dispersion within a projected radius R for a given stellar mass ; then by averaging over all stellar masses, using as statistical weight the total luminosity $\Lambda_{\lambda}^{\delta\lambda}(m)$ of mass component m over a wavelength interval $\delta\lambda$ centered on λ . The mean square velocity of stars of mass m , $\overline{\sigma_m^2}$, is computed from their surface density Σ_m

$$\overline{\sigma_m^2}(R) = \int_0^R \sigma_m^2(r) \Sigma_m(r) \pi dr^2 \Big/ \int_0^R \Sigma_m(r) \pi dr^2$$

The dispersions $\overline{\sigma_m^2}$ are summed over all masses using the luminosity as statistical weight,

$$\langle \sigma_{\lambda}^2 \rangle = \int_{m_0}^{m_1} \overline{\sigma_m^2} \Lambda_{\lambda}^{\delta\lambda}(m[t]) n(m) d \ln m \Big/ \mathcal{W}_{\lambda} \quad (4)$$

where $n(m)$ is the number of stars in the logarithmic mass interval $\ln m, \ln m + \delta \ln m$. The normalisation constant $\mathcal{W}_\lambda$ is the total light flux at wavelength λ from all stars at time t . Note that $n(m)$ in (4) refers to the IMF. All time-dependencies are encapsulated in $\Lambda_\lambda^{\delta\lambda}(m[t])$ and $\overline{\sigma_m^2}$.

For the mass range $1 - 60 M_\odot$ (up to O-stars), the single-star IMF is well represented by a Salpeter (1955) power-law, $n(m)d \ln m \propto m^{-2.35} dm$. We set the mass range $\{m_0, m_1\}$ to cover two decades, from 0.15 to $17 M_\odot$ (spectral type M5 to B5). Note that solar neighbourhood data favour an IMF that flattens out below $1 M_\odot$ (Scalo 1986; Kroupa et al. 1993), so our choice of an Salpeter IMF may seem artificial. However for the chosen mass range, we compute $\langle m \rangle / m_1 \simeq 0.025$, a ratio nearly identical to that obtained from the Galactic-field IMF. Thus our choice of mass function and mass range yields a conservative mass-segregation timescale compared with other IMF’s when the latter are extended beyond $60 M_\odot$.

4 Results

The runtime of our calculations was set to 50 Myrs. All models were scaled to $N = 500,000$ member stars and a total mass of $M \simeq 2 \cdot 10^5 M_\odot$. We vary the dynamical evolution time between runs by constructing models with different sizes and central densities, while keeping the total mass, and N , constant. Mengel et al. (2002) fit the light profile of young Antennae clusters with King models of dimension-less parameter $\Psi/\sigma^2 = 6$ to 9. We first setup three models with identical $\Psi/\sigma^2 = 6$ King parameter but each with different half-light radius (Table 1). The model labeled ‘M82-F’ has a mean surface density $\approx 1.5 \cdot 10^4 M_\odot/pc^2$ or half the value we derive for that cluster from the data of Smith & Gallagher (2001). The densest model is labeled ‘R136’, in reference to the 30 Doradus cluster (central volume density $\sim 10^6 M_\odot/pc^3$; Brandl et al. 1996). Finally, a third low-density model was evolved for comparison (labeled ‘Low’).

We start with the densest model ‘R136’ for which we compute the shortest relaxation time t_{rh} . Since M, r_{ph} and $\langle \sigma_\lambda^2 \rangle$ are all known from the simulations, we solve for η directly from (1). The run of η in time is displayed on Fig. 1(a). Three curves are shown, corresponding to bolometric light and filters centered on the CO continuum and the CaII triplet. The rapid increase of η from an initial value ≈ 8.2 is striking. After 15 Myrs of evolution, η has more than doubled. The projected half-light radius r_{ph} is displayed alongside η . In all the cases r_{ph} decreases steadily in time, a direct result of the migration of massive stars toward the centre. The general trend and quantities are not sensitive to the waveband adopted, for both η and r_{ph} . Note the slow but systematic rise of η after ≈ 35 Myrs, when $\eta > 20$ (factor

> 2.5 from its initial value; Fig. 1[a]).

The LOSVD changes relatively little over time in comparison: we measure a monotonic decrease of $\langle \sigma_\lambda \rangle$ from ≈ 10.4 to 9.3 km/s (or, -10.6%) for the system as a whole, although for individual components evolution was more significant: down $\approx 30\%$ for the most massive stars, while the lightest component enjoys an increase of a comparable magnitude. These effects can all be traced back to the dynamical mass segregation.

The time-evolution of η for all three models is displayed on Fig. 1(b); we have plotted only the results for the bolometric light for clarity. Both η and r_{ph} for the ‘Low’ model stay essentially constant throughout. However the case ‘M82-F’ (initial surface density $> 10M_\odot/\text{pc}^2$) shows unmistakable evolution, suggesting that for clusters of such or higher mean surface density we may no longer presume a time-independent mass-to-light parameter η .

5 Discussion and future work

When the projected density of massive star clusters exceeds a few $\times 10^4 M_\odot/\text{pc}^2$, the mass-to-light conversion factor η used to derive the mass in equation (1) may increase by a factor as large as 3 over time-scales of a few $\times 10^7$ years (cf. Fig. 1[b] and Table 1). Clusters with low surface density have long mass-segregation time-scales and constant η . When we compare clusters with similar mean surface density $\approx 4 \times 10^4 M_\odot/\text{pc}^2$ but different central density initially, the results indicate that the increase in η is an increasing function of the initial central density (Table 1). Thus the drift of heavy stars toward the centre can considerably bias mass estimates of clusters centrally peaked at birth. The resolved cluster R136 (NGC 2070) in the 30 Doradus complex is a case in point: its central volume density may be as high as $\sim 10^6 M_\odot/\text{pc}^3$ with an estimated spherical half-mass radius ~ 1 pc (Brandl et al. 1996). Several clusters in the Antennæ or M82 have estimated masses $\sim 10^6 M_\odot$ and a projected half-light radius $r_{ph} \sim$ a few pc’s. Smith & Gallagher (2001) derive $M = 1.2 \times 10^6 M_\odot$ and $r_{ph} \approx 2.8$ pc for the cluster M82-F. The mean surface density of this cluster $\sim 2.4 \times 10^4 M_\odot/\text{pc}^2$ inside r_{ph} , comparable to our model ‘M82-F’; for comparison, our model ‘R136’ had a mean density twice as high. All calculations were done for a membership of $N = 500,000$, a factor 2 lower than rich starburst clusters. This may serve to accelerate evolution compared with actual massive clusters (cf. Eq. [2]). We recall that our choice of an upper mass $m_1 < 20 M_\odot$ yields a mass segregation timescale which is a factor of 3 longer than if we had included O-stars. A re-run of the case ‘M82-F’ with a mass spectrum widened to $50 M_\odot$ yielded a rapid increase of η over a shorter, 10 Myrs-timescale (open dots, Fig. 1[b]). Therefore the evolution of η through mass segregation of stars is a sensitive function of both the IMF and the cluster surface density profiles. We mention that the cluster M82-F

has an age of $\simeq 60 \pm 20$ Myrs, by which time the increase of η is maximum. This larger value of η would go some way toward solving the apparent over-luminosity of young M82 clusters with respect to a standard IMF. Recent work on M82-F has highlighted the possibility of strong mass segregation in that cluster (McCrady et al. 2004).

The strong evolution of η for model ‘R136’ (Fig. 1[a]) suggested to us to compare functional fits to the total surface density and luminosity profiles. We performed least-square fits (Press et al. 1992) using King profiles on the simulation grid out to $5r_{ph}$. Doing this for model ‘R136’ after 35 Myrs of evolution we found a best fit to the radial luminosity profile (cf. §3) that gave a King parameter $\Psi/\sigma^2 \approx 7.6$ (compared with $\simeq 6$ for the run of density). The concentration parameter $c = \log |r_t/r_o|$ increases by ≈ 0.45 as Ψ/σ^2 runs from 6 to 7.6. Since the truncation radius r_t remains constant, the central region seemingly shrinks by a factor ≈ 2.8 compared with the surface density. Table 1 gives values of $\langle \eta \rangle$ and $\langle r_{ph} \rangle$ averaged over the last 25 Myrs of evolution for this and two other models, which also show increased concentration. The effect can be cast in the context of a whole population of clusters as detected in interacting galaxies such as the Antennæ (NGC 4038/9) or M82. Our modelling indicates that the clusters should be yet more massive than estimated until now, and less concentrated. Therefore their binding energy per unit mass will be less than deduced from profiling the light. However it has not escaped our attention that fitting the light flux would require a convolution with the point-spread function of an instrument to allow for direct comparisons with observations. More detailed models will be presented in a follow-up study (Fleck et al., in preparation).

Acknowledgments

We thank the anonymous referee for a first-class report. C.M.B. and A.L. thank A.S.S.N.A. and the P.N.G. (France) for financial support. D.C.H. and S.D. warmly thank Observatoire de Strasbourg for hospitality during the course of this research.

REFERENCES

- Binney, J. & Tremaine, S. 1987, *Galactic Dynamics*, Princeton: University of Princeton Press, 733 ff
- Brandl,B., Sams,B.J., Bertoldi,F. et al. 1996, *ApJ* 466, 254
- Eggleton, P.P., Tout, C.A. & Fitchett, M.J. 1989, *ApJ* 347, 998
- Farouki, R.T. & Salpeter, E. E. 1982, *ApJ* 253, 512
- Giersz, M. & Spurzem, R. 1994, *MNRAS* 269, 241
- Heggie, D.C. & Ramamani, N. 1988, *MNRAS* 237, 757
- Hurley,J.R., Pols,O.R. & Tout,C.A. 2000, *MNRAS* 315, 543
- King, I.R. 1966, *AJ* 71, 64
- Kroupa, P. 2002, *Science* 295, 82
- Kroupa, P., Tout, C.A. & Gilmore, G. 1993, *MNRAS* 262, 545
- Larson, R. B. 1970, *MNRAS* 147, 323
- Lejeune, T., Cuisinier,F. & Buser,R. 1998, *A&AS* 130, 65
- Louis, P.D. & Spurzem, R. 1991, *MNRAS* 251, 408
- Lynden-Bell, D. & Eggleton, P.P. 1980, *MNRAS* 191, 483
- Maraston,C., Bastian,N., Saglia,R.P. et al. 2004, *A&A* 416, 467
- McCrady, N., Gilbert, A.M. & Graham, J.R. 2003, *ApJ* 596, 240
- McCrady, N., Graham, J.R. & Vacca, W.D. 2004, *astro-ph/0411256*
- Mengel, S., Lehnert, M.D., Thatte, N. et al. 2002, *A&A* 383, 137
- Meurer,G.R.; Heckman, T.M.; Leitherer,C. et al. 1995, *AJ* 110, 2665
- Meylan, G. & Heggie, D.C. 1997, *A&A Reviews* 8, 1
- Pols O.R., et al. 1998, *MNRAS* 298, 525
- Press, W.H. et al. 1992, *Numerical Recipes*, Cambridge: University Press, 963 pp

- Salpeter, E.E. 1955, ApJ 121, 161
- Scalo, J.M. 1986, Fund. Cosmic Physics, 11, 1
- Schweizer, F. 1986, Science 231, 227
- Smith, L.J. & Gallagher, J. S. III 2001, MNRAS 326, 1027
- Spitzer, L. jr 1987, The Dynamics of Star Clusters, Princeton: University Press
- Spitzer, L. Jr & Shull, J.M. 1975, ApJ 201, 773
- Spurzem, R. & Takahashi, K. 1995, MNRAS 272, 772
- Sternberg, A. 1998, ApJ 506, 721
- Suntzeff, N. 1993, in The Globular Clusters – Galaxy Connection, ASP Conf. Series Vol. 48, ed. G.H. Smith & J.P. Brodie (San Francisco: PASP), 167




Fig. 1.— Parameter η and half-light radius r_{ph} versus time. The initial models were unsegregated King $\Psi/\sigma^2 = 6$ models. a) results using filters at three different wavebands (see text for details); b) results for models with different initial surface density (cf. Table 1). When the mass range is increased from $17 M_{\odot}$ to $50 M_{\odot}$ η rises sharply over the first 10 Myrs (open circles, case ‘M82-F’); thereafter it rejoins the curve displayed.

Table 1: Parameters of the models. All models have a total mass $M = 2 \times 10^5 M_\odot$. We set $R_h = R_g/2$ and $\rho = \bar{\rho}$ in (3) to average t_{ms} . The subscript 0 denotes $t = 0$; hats denote time-averages over the last 25 Myrs of evolution. The last column gives the increase of concentration parameter δc for a few cases (§§5.1, 5.2).

Name	Model Ψ/σ^2	t_{ms} Myrs	$r_{ph,0}$ pc	$\widehat{r_{ph}}$ pc	$\Sigma_0(0)$ $10^5 M_\odot/pc^2$	$\overline{\Sigma_0}(r_{ph})$ $10^5 M_\odot/pc^2$	η_o	$\widehat{\eta}$	$\widehat{\delta c}$
‘R136’	6	9.3	1.01	0.51 ± 0.03	0.885	0.417	8.15	18.5 ± 1	0.29
‘M82-F’	6	15.9	1.46	1.02 ± 0.06	0.474	0.224	8.15	12.6 ± 0.7	
Low	6	47.9	3.05	2.84 ± 0.05	0.109	0.051	8.15	8.54 ± 0.06	
‘Plummer’	4.5	7.5	1.05	0.62 ± 0.03	0.606	0.410	8.20	16 ± 1	0.23
	9	25.7	0.95	0.49 ± 0.02	5.232	0.440	7.76	19 ± 4	0.29